



# Mg/capita Consumption of Fentanyl, Burkina Faso, 1980-2007




Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

# Mg/capita Consumption of Hydromorphone, Burkina Faso, 1980-2007




Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

# Mg/capita Consumption of Methadone, Burkina Faso, 1980-2007




Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

# Mg/capita Consumption of Morphine, Burkina Faso, 1980-2007




Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

# Mg/capita Consumption of Oxycodone, Burkina Faso, 1980-2007



Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

# Mg/capita Consumption of Pethidine, Burkina Faso, 1980-2007



Sources: International Narcotics Control Board; United Nations population data  
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009