



Mg/capita Consumption of Fentanyl, Burundi, 1980-2007




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Hydromorphone, Burundi, 1980-2007




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Methadone, Burundi, 1980-2007




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Morphine, Burundi, 1980-2007




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Oxycodone, Burundi, 1980-2007



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Pethidine, Burundi, 1980-2007



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008