



Mg/capita Consumption of Fentanyl, Belgium, 1980-2006




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Hydromorphone, Belgium, 1980-2006




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Methadone, Belgium, 1980-2006




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Morphine, Belgium, 1980-2006




Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Oxycodone, Belgium, 1980-2006



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Mg/capita Consumption of Pethidine, Belgium, 1980-2006



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008