


WPRO Consumption of Pethidine, 2007



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

WPRO Regional Pethidine Consumption, 2007: (mg/capita)

Note: Countries not listed did not report pethidine consumption to the INCB for 2007

| | | |
|----|---------------------|--------|
| 1 | Cook Islands | 8.6667 |
| 2 | New Zealand | 7.9395 |
| 3 | Australia | 5.1665 |
| 4 | Hong Kong SAR | 2.8665 |
| 5 | Tonga | 2.6900 |
| 6 | Republic of Palau | 2.5909 |
| 7 | Brunei Darussalam | 2.5775 |
| 8 | Singapore | 2.5044 |
| 9 | Republic of Korea | 2.4347 |
| 10 | Macao | 2.1707 |
| 11 | Papua New Guinea | 2.0505 |
| 12 | Samoa | 1.5200 |
| 13 | Marshall Islands | 1.3427 |
| 14 | China | 1.1906 |
| 15 | Viet Nam | 0.9350 |
| 16 | Lao Peop. Dem. Rep. | 0.6040 |
| 17 | Japan | 0.3986 |
| 18 | French Polynesia | 0.2235 |
| 19 | Philippines | 0.1460 |
| 20 | Cambodia | 0.0001 |



Pain & Policy Studies Group
WHO Collaborating Center for Policy and
Communications in Cancer Care
University of Wisconsin Carbone Cancer Center
School of Medicine and Public Health