


2006 Global Consumption of Hydromorphone



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2008

Global Hydromorphone Consumption, 2006: (mg/capita)

Note: Countries not listed did not report hydromorphone consumption to the INCB for 2006

1	Canada	17.1220
2	Austria	4.4013
3	United States of America	3.5971
4	Germany	2.8407
5	Belgium	1.5651
6	Sweden	1.4449
7	Ireland	0.8168
8	Switzerland	0.6876
9	France	0.5027
10	Slovenia	0.4680
11	Australia	0.4032
12	Luxembourg	0.3920
13	Czech Republic	0.3176
14	Slovakia	0.2250
15	Iceland	0.2100
16	Denmark	0.2063
17	Norway	0.1791
18	Netherlands	0.1785
19	Hungary	0.1636
20	United Kingdom	0.1304
21	Finland	0.1279
22	New Caledonia	0.0783
23	Israel	0.0166
24	French Polynesia	0.0157
25	Saudi Arabia	0.0142
26	Colombia	0.0115
27	Croatia	0.0059
28	Mexico	0.0030
29	Republic of Korea	0.0020
30	South Africa	0.0002



Pain & Policy Studies Group
WHO Collaborating Center for Policy and Communications in
Cancer Care
University of Wisconsin- School of Medicine and Public Health
Paul P. Carbone Comprehensive Cancer Center