


2007 Global Consumption of Hydromorphone



Sources: International Narcotics Control Board; United Nations population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2009

Global Hydromorphone Consumption, 2007: (mg/capita)

Note: Countries not listed did not report hydromorphone consumption to the INCB for 2007

1	Canada	18.5850
2	Austria	5.5648
3	United States of America	3.8594
4	Germany	3.6795
5	Sweden	1.8854
6	Slovenia	1.2990
7	Belgium	1.0901
8	Switzerland	0.9281
9	Ireland	0.8361
10	Czech Republic	0.6181
11	Luxembourg	0.5840
12	Iceland	0.4367
13	Denmark	0.4304
14	Australia	0.4261
15	France	0.3871
16	Slovakia	0.2550
17	Hungary	0.1937
18	Netherlands	0.1833
19	Finland	0.1706
20	French Polynesia	0.1451
21	Norway	0.1126
22	New Caledonia	0.1000
23	United Kingdom	0.0950
24	Colombia	0.0474
25	Italy	0.0373
26	Croatia	0.0309
27	Republic of Korea	0.0164
28	Israel	0.0116
29	Saudi Arabia	0.0075
30	United Arab Emirates	0.0073
31	Mexico	0.0061
32	Poland	0.0021
33	Serbia	0.0004
34	Greece	0.0004



Pain & Policy Studies Group
WHO Collaborating Center for Policy and
Communications in Cancer Care
University of Wisconsin Carbone Cancer Center
School of Medicine and Public Health