

Making Opioid Pain Medications Available to Patients is an Essential Part of Cancer Control

WHO Consultation on Strategies to Improve and Strengthen Cancer Control Programmes in Europe

25-28 November 2003



World Health Organization Collaborating Center for
Policy and Communications in Cancer Care
Pain & Policy Studies Group
University of Wisconsin Comprehensive Cancer Center
www.medsch.wisc.edu/painpolicy

Selected Bibliography

Blengini C, Joranson DE, Ryan KM. Italy reforms national policy for cancer pain relief and opioids. *European Journal of Cancer Care*. 2003;12(1):28-34. <http://www.medsch.wisc.edu/painpolicy/publicat/03ejcc/index.htm>.

International Narcotics Control Board. *Report of the International Narcotics Control Board for 1995: Availability of opiates for medical needs*. New York, NY: United Nations; 1996. http://www.incb.org/e/ind_ar.htm.

Joranson DE. Improving availability of opioid pain medications: Testing the principle of balance in Latin America. *Innovations in End-of-Life Care*. 2003;5(1). <http://www2.edc.org/lastacts/archives/archivesJan03/featureinn.asp>.

Pain & Policy Studies Group. *Improving cancer pain relief in the world: Report for 2002*. Madison, Wisconsin, USA: University of Wisconsin Pain & Policy Studies Group/WHO Collaborating Center for Policy and Communications in Cancer Care; 2003. <http://www.medsch.wisc.edu/painpolicy/publicat/annrepts.htm>.

Pain & Policy Studies Group. Opioid consumption statistics in 6 Eastern European countries and the region. http://www.medsch.wisc.edu/painpolicy/internat/E.Europe/Eastern_Europe.htm

United Nations. *Competent national authorities under the international drug control treaties, 2000*. New York, NY: United Nations; 2001.

World Health Organization. *Cancer pain relief with a guide to opioid availability (2nd edition)*. Geneva, Switzerland: World Health Organization; 1996. <http://whqlibdoc.who.int/publications/9241544821.pdf>.


World Health Organization. *Cancer pain relief and palliative care in children*. Geneva, Switzerland: World Health Organization; 1998. <http://whqlibdoc.who.int/publications/9241545127.pdf>

World Health Organization. *Achieving balance in national opioids control policy: Guidelines for assessment*. Geneva, Switzerland: World Health Organization; 2000. <http://www.medsch.wisc.edu/painpolicy/publicat/00whoabi/00whoabi.htm>.

World Health Organization. *National cancer control programmes: Policies and managerial guidelines (2nd edition)*. Geneva, Switzerland: World Health Organization; 2002. <http://www.who.int/cancer/nccp/nccp/en/>.

World Health Organization. *World cancer report*. Lyon, France: IARC Press; 2003. <http://bookorders.who.int/newaccess/anglais/detart1.jsp?sesslan=1&codlan=1&codcol=76&codcch=0016#>


Global Consumption of Morphine, Oxycodone & Pethidine 1985-2001



Source: International Narcotics Control Board

By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003


Global Consumption of Fentanyl 1985-2001



Source: International Narcotics Control Board


By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003

Consumption of Morphine in Europe mg/capita, 2001




Source: International Narcotics Control Board; United Nations Demographic Yearbook
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003

Consumption of Fentanyl in Europe mg/capita, 2001




Source: International Narcotics Control Board; United Nations Demographic Yearbook
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003

Consumption of Oxycodone in Europe mg/capita, 2001



Source: International Narcotics Control Board; United Nations Demographic Yearbook
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003

Consumption of Pethidine in Europe mg/capita, 2001



Source: International Narcotics Control Board; United Nations Demographic Yearbook
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2003