


Burundi

Opioid Consumption in Morphine Equivalence (ME), mg per person



Sources: International Narcotics Control Board; World Health Organization population data
 By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2015

Burundi

Opioid Consumption in Morphine Equivalence (ME), mg per person

Note: Blank cells indicate no consumption data

Year	Methadone ME	Morphine	Pethidine ME	Oxycodone ME	Fentanyl ME	Hydromorphone ME	Total ME
1980							0.0000
1981							0.0000
1982							0.0000
1983			0.1439				0.1439
1984		0.0109	0.0291				0.0400
1985		0.0011	0.0161				0.0171
1986					0.1716		0.1716
1987							0.0000
1988							0.0000
1989							0.0000
1990							0.0000
1991					0.0593		0.0593
1992							0.0000
1993							0.0000
1994							0.0000
1995							0.0000
1996							0.0000
1997							0.0000
1998		0.0283	0.0035		0.0011		0.0328
1999							0.0000
2000		0.0008	0.0046		0.0092		0.0145
2001			0.0258		0.0081		0.0339
2002		0.0030	0.0155		0.0211		0.0396
2003		0.0951	0.0320		0.0079		0.1350
2004		0.0080	0.0142		0.0054		0.0275
2005		0.0025	0.0128		0.0017		0.0170
2006		0.0019	0.0288		0.0067		0.0374
2007		0.0118	0.0769				0.0887
2008		0.0050			0.0081		0.0131
2009							0.0000
2010							0.0000
2011							0.0000
2012							0.0000
2013							0.0000
2014							0.0000
2015		0.0310	0.0022		0.0587		0.0920

Data sources:

Consumption data - International Narcotics Control Board
 Population - United Nations World Population Prospects, 2013 Revision
 ME conversion factors - WHOCC Centre for Drug Statistics Methodology

Pain & Policy Studies Group
 University of Wisconsin
 Carbone Cancer Center
 WHO Collaborating Center