


United Kingdom

Opioid Consumption in Morphine Equivalence (ME), mg per person



Sources: International Narcotics Control Board; World Health Organization population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2015

United Kingdom

Opioid Consumption in Morphine Equivalence (ME), mg per person

Note: Blank cells indicate no consumption data

Year	Methadone ME	Morphine	Pethidine ME	Oxycodone ME	Fentanyl ME	Hydromorphone ME	Total ME
1980	3.4226	3.5066	2.0417	0.1249	0.2204		9.3163
1981	3.6557	4.3176	2.0380	0.1805	0.2957		10.4874
1982	4.2864	3.8038	1.1509	0.2072	0.0562		9.5045
1983	4.4741	2.7999	1.7188	0.3725	0.3904		9.7557
1984	4.4669	8.1487	1.6636	0.2803	0.3067	0.0005	14.8667
1985	4.6254	7.5350	1.7260	0.3941	0.6130	0.0026	14.8960
1986	4.6409	10.3601	1.6023	0.2025	0.3884	0.0123	17.2065
1987	4.9744	12.9366	1.6115	0.0952	0.8098	0.1101	20.5376
1988	6.5199	14.4090	1.5749	0.0811	0.5555	0.1944	23.3347
1989	8.5978	14.7685	1.7709	0.1759	0.1322	0.2655	25.7108
1990	12.9874	16.3012	2.2185	0.0747	0.4097	0.7912	32.7827
1991	14.4834	20.0820	1.8034	0.0325	0.4916	0.0020	36.8948
1992	23.3395	14.8254	6.6581	0.0252	0.5876	0.4372	45.8730
1993	28.5102	37.7348	1.5478	0.1891	0.5785	0.2176	68.7780
1994	35.5406	36.3043	3.1252	0.0390	0.9647	0.6928	76.6665
1995	14.1166	25.6217	3.1874	0.3900	2.2039	0.4116	45.9313
1996	36.7759	18.8262	2.1003	0.0239	4.0842	0.0441	61.8547
1997	39.0284	21.9802	2.1910	0.0381	15.3403	0.1103	78.6882
1998	51.2442	20.4646	1.9586	0.0380	18.1960	0.4037	92.3050
1999	43.9519	19.7325	1.6386	0.1375	24.5030	0.6302	90.5937
2000	33.9434	19.1616	1.6897	1.2613	21.7426	0.3748	78.1733
2001	38.5583	18.7539	1.2107	1.7975	11.7732	0.6263	72.7200
2002	53.9902	18.7036	1.2665	3.2029	18.7940	0.5255	96.4828
2003	55.7459	19.3842	1.1855	5.6019	19.4554	0.4407	101.8136
2004	62.0357	19.1548	0.4558	8.1102	27.1035	0.3802	117.2402
2005	80.0684	28.5573	1.3270	11.2058	46.5892	0.5784	168.3261
2006	71.1942	28.2392	0.8267	9.1830	36.2928	0.6520	146.3879
2007	49.1809	21.1144	0.7073	4.7920	13.3278	0.4752	89.5977
2008	159.0956	41.2246	1.2018	39.8061	55.5999	0.4401	297.3680
2009	197.7273	56.8781	1.6344	52.7571		2.9612	311.9581
2010	193.4361	56.4189	0.8576	21.2896	551.5089	0.7863	824.2973
2011	157.1367	48.0638	1.6262	17.6859	28.0361	0.4005	252.9492
2012	90.6943	39.2065	0.5662	31.0172	83.2591	0.6771	245.4204
2013	128.5667	33.4069	0.4581	18.3692	57.7923	2.8764	241.4695
2014	161.0574	48.7475	0.4347	121.7336	91.8129	0.6242	424.4104
2015	124.2615	22.8571	0.6207	26.5038	106.1583	0.7334	281.1348

Data sources:

Consumption data - International Narcotics Control Board
 Population - United Nations World Population Prospects, 2013 Revision
 ME conversion factors - WHOCC Centre for Drug Statistics Methodology

Pain & Policy Studies Group
 University of Wisconsin
 Carbone Cancer Center
 WHO Collaborating Center