


Burkina Faso

Opioid Consumption in Morphine Equivalence (ME) minus Methadone, mg per person



Sources: International Narcotics Control Board; World Health Organization population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2015

Burkina Faso

Opioid Consumption in Morphine Equivalence (ME) minus Methadone, mg per person

Note: Blank cells indicate no consumption data

Year	Morphine	Pethidine ME	Oxycodone ME	Fentanyl ME	Hydromorphone ME	Total ME
1980						0.0000
1981						0.0000
1982						0.0000
1983	0.0001	0.0047				0.0048
1984	0.0001	0.0094				0.0096
1985		0.0017				0.0017
1986	0.0001	0.0057				0.0058
1987	0.0001	0.0055				0.0056
1988	0.0008	0.0051				0.0059
1989		0.0076		0.0095		0.0171
1990	0.0021	0.0126				0.0147
1991	0.0004	0.0050				0.0054
1992	0.0001	0.0025				0.0026
1993	0.0005	0.0052				0.0057
1994		0.0034				0.0034
1995		0.0061		0.0127		0.0189
1996		0.0061		0.0096		0.0157
1997						0.0000
1998						0.0000
1999						0.0000
2000		0.0113		0.0101		0.0215
2001						0.0000
2002	0.0006			0.0055		0.0061
2003		0.0018		0.0209		0.0228
2004	0.0016			0.0245		0.0261
2005	0.0007			0.0201		0.0208
2006	0.0017			0.0124		0.0141
2007	0.0007			0.0043		0.0050
2008	0.0026	0.0008		0.0044		0.0077
2009	0.0046	0.0004		0.0164		0.0214
2010						0.0000
2011	0.0128			0.0175		0.0303
2012	0.0099	0.0002		0.0052		0.0153
2013						0.0000
2014	0.0139			0.0140		0.0279
2015	0.0190			0.0157		0.0347

Data sources:

Consumption data - International Narcotics Control Board

Population - United Nations World Population Prospects, 2013 Revision

ME conversion factors - WHOCC Centre for Drug Statistics Methodology

Pain & Policy Studies Group

University of Wisconsin

Carbone Cancer Center

WHO Collaborating Center