


Anguilla

Opioid Consumption in Morphine Equivalence (ME), mg per person



Sources: International Narcotics Control Board; World Health Organization population data
By: Pain & Policy Studies Group, University of Wisconsin/WHO Collaborating Center, 2015

Anguilla

Opioid Consumption in Morphine Equivalence (ME), mg per person

Note: Blank cells indicate no consumption data

Year	Methadone ME	Morphine	Pethidine ME	Oxycodone ME	Fentanyl ME	Hydromorphone ME	Total ME
1980							0.0000
1981			1.2500				1.2571
1982							0.0000
1983		1.7143	2.3929				4.1000
1984		0.1429	1.8571				2.0000
1985		0.1429	1.6071				1.7571
1986			1.5000				1.5000
1987		0.1429	2.0714				2.2143
1988		0.1429	1.1429				1.2857
1989							0.0000
1990							0.0000
1991							0.0000
1992			2.2222				2.2222
1993							0.0000
1994							0.0000
1995							0.0000
1996							0.0000
1997							0.0000
1998		0.0833	0.7083				0.7917
1999							0.0000
2000			0.5455		0.0076		0.5545
2001			0.7955		0.0455		0.8364
2002		0.0833	1.2500		0.1389		1.4750
2003							0.0000
2004		0.0833	0.7292		0.0417		0.8500
2005			0.0385		0.3846		0.4231
2006							0.0000
2007							0.0000
2008		0.3333	0.9333				1.2667
2009			0.1333		0.0389		0.1733
2010		0.5333	0.6667		0.0389		1.2400
2011		0.5333	1.0500		0.1611		1.7467
2012							0.0000
2013		0.0635	0.4602				0.5269
2014		0.4973	0.1088		0.0311		0.6341
2015		0.3333	0.5833		0.0889		1.0067

Data sources:

Consumption data - International Narcotics Control Board

Population - United Nations World Population Prospects, 2013 Revision

ME conversion factors - WHOCC Centre for Drug Statistics Methodology

Pain & Policy Studies Group

University of Wisconsin

Carbone Cancer Center

WHO Collaborating Center